

Regional Variations in FUV Lunar Signatures: **Swirls**

Amanda R. Hendrix

T. K. Greathouse, K. D. Retherford,

K. E. Mandt, G. R. Gladstone,


D. E. Kaufmann, D. M. Hurley, P. D. Feldman,

W. R. Pryor, M. A. Bullock, S. A. Stern



Swirls: a quick overview

- Swirls are associated with areas of magnetized crust
- Swirls exhibit spectral characteristics (VNIR) similar to immature material
- Possible causes:
 - Stand-off of solar wind (e.g. Glotch et al., 2015)
 - Inhibiting nanophase iron production (except via micrometeoroid bombardment)
 - Could also enhance weathering in surrounding regions (Kramer et al., 2011)
 - Dust transport caused by electric fields (e.g. Garrick-Bethell et al., 2011)
 - Accumulating fine-grained feldspathic particles in these regions



LAMP:

FUV Spectrograph

Mass: 6.08 kg

Power: 4.8 W



λ Range: 57-196 nm

FOV: $0.3^\circ \times 6.0^\circ$

Filled-Slit λ Range: <4 nm

LRO LAMP

Lyman Alpha Mapping Project



FUV: Surface scattering regime


$$R \propto \frac{(n-1)^2 + k^2}{(n+1)^2 + k^2}$$

FUV: ~ 100 - 200 nm


NUV: ~ 200 - 350 nm



Swirls



from Kramer et al., 2011




Here we focus on Reiner & Gerasimovich



from Kramer et al., 2011

UV ratio images

- For these ratio (180/150 nm) images, generally
 - Highlands: higher ratio values (red, less blue)
 - Maria: lower ratio values (blue)
 - Weathered regions: lower ratio values
 - Less-weathered regions: higher ratio values (less blue)




Reiner Gamma




Ratio ($\sim 180\text{-}190\text{ nm}/150\text{ nm}$); with
phase correction

summing 22 months of data with $\beta < 70^\circ$



Reiner Gamma



Reiner Gamma





Reiner Gamma



The visibly-bright swirl region is less blue in FUV than surrounding terrain



Gerasimovich swirl





Ratio ($\sim 180\text{-}190\text{ nm}/150\text{ nm}$);
with phase correction

Summing 34 months of data with $\text{beta} < 80^\circ$

Gerasimovich swirl




Gerasimovich swirl



The visibly-bright swirl region is significantly redder in FUV than surrounding terrain

Bluing could be due to increased pyroxene and/or ilmenite.
Gerasimovich swirls have spectral signature similar to anorthite.




data from Wagner et al., 1987



UV Weathering Effect (#1)

- Lunar space weathering is (**in the VNIR**) characterized by spectral reddening, darkening and degradation of absorption bands. All 3 effects are related to nanophase iron.
 - In the swirls, the normal weathering pattern is not seen – not much darkening or reddening, though absorption bands are weak.
- **In the NUV & FUV**, lunar soils are bluer than rocks (Wagner et al., 1987; Hendrix & Vilas, 2006)
 - Linking the bluing to weathering

Lunar soils: thick; Lunar rocks: thin/diamonds




data from Wagner et al., 1987



Note that FUV ratios are <1 , even for bright rays


UV Weathering Effect (#2)



Mature regions are bluer than immature regions we also know from lunar samples.


Immature regions are darker than mature regions - attributed to a lack of SMFe and/or a lack of gardening.

Swirls are not just immature, unweathered regions.



Models of a feldspar with increasing abundances of SMFe inclusions
(after Hapke, 2001).

Gerasimovich swirl is similar to a somewhat-weathered feldspar.



Summary

- Swirls are detected in LAMP FUV data
 - spectrally red compared to surrounding terrains
 - swirls are weathered to a certain extent: spectrally different from immature regions (immature soils are darker)
 - however, swirls do not show FUV spectral evidence of being mature (mature soils are blue)
- Immature soils (e.g. bright rays) are FUV dark due to a relative lack of nanophase iron (and/or a lack of gardening) and spectrally (a little) blue, likely due to pyroxene content.
- Mature soils are relatively bright due to the nanophase iron content (and/or gardening) and are bluer than immature soils, perhaps due to increased ilmenite and/or iron, especially in weather-produced grain rims
- Swirls are relatively bright like mature soils are spectrally red, not blue
 - spectrally consistent with moderately weathered feldspar
 - Consistent with the dust transport/compositional sorting model (Garrick-Bethell et al., 2011)
- Overall, we find these characteristics are consistent with a model where the magnetically anomalous swirls stand off much of the solar wind (inhibiting a bluing of the spectrum)
 - micrometeoroid bombardment acts to weather swirls
 - Gives us a clue to distinguish solar wind effects from micrometeoroid bombardment effects
 - Micrometeoroid bombardment brightens as it gardens and/or adds SMFe
 - these FUV lunar weathering effects will be described more fully in a future paper